



Apr 4th, 1:30 PM - 1:45 PM

Assessing harmful algal bloom risk in Puget Sound: a coupled modeling-data analysis approach

D. L. Woodruff

Pacific Northwest National Laboratory (U.S.), dana.woodruff@pnnl.gov

Taiping Wang

Pacific Northwest National Laboratory (U.S.), taiping.wang@pnnl.gov

Stephanie K. Moore

National Marine Fisheries Service (U.S.), stephanie.moore@noaa.gov

Zhaoqing Yang

Pacific Northwest National Laboratory (U.S.), zhaoqing.yang@pnnl.gov

Ning Sun

Pacific Northwest National Laboratory (U.S.), Ning.Sun@pnnl.gov

See next page for additional authors

Follow this and additional works at: <https://cedar.wvu.edu/ssec>



Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Woodruff, D. L.; Wang, Taiping; Moore, Stephanie K.; Yang, Zhaoqing; Sun, Ning; Borchert, Jerry; Coyne, Audrey; Mauger, Guillaume; and Cullinan, Valerie, "Assessing harmful algal bloom risk in Puget Sound: a coupled modeling-data analysis approach" (2018). *Salish Sea Ecosystem Conference*. 14.
<https://cedar.wvu.edu/ssec/2018ssec/allsessions/14>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wvu.edu.

Speaker

D. L. Woodruff, Taiping Wang, Stephanie K. Moore, Zhaoqing Yang, Ning Sun, Jerry Borchert, Audrey Coyne, Guillaume Mauger, and Valerie Cullinan



Assessing harmful algal bloom risk in Puget Sound – a coupled modeling-data analysis approach

**Dana Woodruff, Taiping Wang, Zhaoqing Yang, Val Cullinan,
Ning Sun, Rebecka Bence**
(Pacific Northwest National Laboratory)

Stephanie Moore
(NOAA NW Fisheries Science Center)

Guillaume Mauger
(Univ. of Wash. Climate Impacts Group)

Jerry Borchert, Audrey Coyne
(WA Dept. of Health)

Salish Sea Ecosystem Conference April 4th, 2018



Background and Goals

- ▶ Projected increase of HAB threats to shellfish harvest and associated economic impact to coastal communities
- ▶ Need for better forecasting tools for HAB occurrence and biotoxin risk
- ▶ Monitoring resources limited, strategic resource allocations necessary

GOAL:

Better understand HAB risk to support improved management of shellfish monitoring now and in the future



Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by **Battelle** Since 1965


Why Paralytic Shellfish Poisoning (PSP)?

- ▶ *Alexandrium* bloom in Washington's marine waters every year
- ▶ PSP is a serious illness caused by eating contaminated shellfish
- ▶ Over 3,300 shellfish samples are tested annually for PSP
- ▶ Biotoxin monitoring is a coordinated effort between DOH, DFW, DNR, Local Health Departments, Tribes, Industry and Citizen Volunteers
- ▶ Variability in the timing and location of PSP blooms presents planning challenges to managers

Phytoplankton tow 9/26/17
Credit: Penn Cove Shellfish, LLC




Development of PSP Risk Probabilities for Puget Sound



High-resolution Salish Sea Hydrodynamic Model

- ***Inputs to model:***
 - Hydrology and climate forcing from UW DHSVM and WRF
- ***Outputs from model:***
 - Temperature/salinity
 - Light availability
 - Residence time
 - Vertical stratification
- ***Simulation periods:***
 - Recent: 2000-2006
 - Future: mid-century




PSP Shellfish Harvest Closure Probabilities

- ▶ Mussel sentinel sites sampled year round
- ▶ 27 biotoxin closure zones located in 12 coastal counties
- ▶ # of closure days
 - Each site/zone
 - Weekly/biweekly
 - 2000-2006




Regional scale










Closure zone example – Bellingham Bay







Closure zone example – Bellingham Bay












































Model Skill Assessment



Probability of closure





Next steps

- ▶ Assess probability of shellfish closure for future time period
 - Does the pattern of PSP risk change in the future?
 - How will this affect monitoring effort?



Conclusions

- ▶ Application of modeling approach to the scale of management units (biotoxin closure zones) is unique in Puget Sound
- ▶ Predicts seasonality of shellfish closure probabilities reasonably well
- ▶ Interannual variability of closures not well captured at all sites, hence...
- ▶ Not necessarily a good tool for next year predictions, but potentially a good tool for assessing longer-term (decadal scale) changes to the PSP bloom season

Questions?

Contact information:

Dana Woodruff

Ph: 360-681-3608

Dana.Woodruff@pnnl.gov



Acknowledgements: Funding provided by NOAA's Coastal and Ocean Climate Applications Program